



# FINAL JEE-MAIN EXAMINATION - JULY, 2021

# (Held On Thursday 22<sup>nd</sup> July, 2021)

TIME: 3:00 PM to 6:00 PM

## CHEMISTRY

# SECTION-A

The water having more dissolved O<sub>2</sub> is :

 boiling water
 water at 80°C
 polluted water
 water at 4°C

# Official Ans. by NTA (4)

- 2. Which one of the following statements for D.I. Mendeleeff, is **incorrect**?
  - He authored the textbook Principles of Chemistry.
  - (2) At the time, he proposed Periodic Table of elements structure of atom was known.
  - (3) Element with atomic number 101 is named after him.
  - (4) He invented accurate barometer.


#### Official Ans. by NTA (2)

- **3.** Which purification technique is used for high boiling organic liquid compound (decomposes near its boiling point)?
  - (1) Simple distillation
  - (2) Steam distillation
  - (3) Fractional distillation
  - (4) Reduced pressure distillation

## Official Ans. by NTA (4)

**4.** Which of the following compounds will provide a tertiary alcohol on reaction with excess of CH<sub>3</sub>MgBr followed by hydrolysis?

**TEST PAPER WITH ANSWER** 



#### Official Ans. by NTA (1)

Which of the following compounds does not exhibit resonance?

(1) CH<sub>3</sub>CH<sub>2</sub>OCH=CH<sub>2</sub>

(3) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CONH<sub>2</sub>
(4) CH<sub>3</sub>CH<sub>2</sub>CH=CHCH<sub>2</sub>NH<sub>2</sub>
Official Ans. by NTA (4)

5.



Match List-I with List-II			
List-I	List-II		
(Elements)	(Properties)		
(a) Ba	(i) Organic solvent soluble		
	compounds		
(b) Ca	(ii) Outer electronic configuration		
	$6s^2$		
(c) Li	(iii) Oxalate insoluble in water		
(d) Na	(iv) Formation of very strong		
	monoacidic base		
Change the compact enginer from the entire give			

Choose the **correct** answer from the options given below :


(1) (a)-(ii), (b)-(iii), (c)-(i) and (d)-(iv)

(2) (a)-(iv), (b)-(i), (c)-(ii) and (d)-(iii)

(3) (a)-(iii), (b)-(ii), (c)-(iv) and (d)-(i)

(4) (a)-(i), (b)-(iv), (c)-(ii) and (d)-(iii)

Official Ans. by NTA (1)



In the chemical reactions given above A and B respectively are :

(1) H<sub>3</sub>PO<sub>2</sub> and CH<sub>3</sub>CH<sub>2</sub>Cl

(2) CH<sub>3</sub>CH<sub>2</sub>OH and H<sub>3</sub>PO<sub>2</sub>

(3) H<sub>3</sub>PO<sub>2</sub> and CH<sub>3</sub>CH<sub>2</sub>OH

(4)  $CH_3CH_2Cl$  and  $H_3PO_2$ 

## Official Ans. by NTA (1)

8. Isotope(s) of hydrogen which emits low energy  $\beta^-$  particles with  $t_{\frac{1}{2}}$  value > 12 years is/are

(1) Protium

- (2) Tritium
- (3) Deuterium
- (4) Deuterium and Tritium

Official Ans. by NTA (2)

9. Match List-I with List-II :

List-I		List-II
(Species)		(Hybrid Orbitals)
(a) SF <sub>4</sub>	(i)	$sp^{3}d^{2}$
(b) IF <sub>5</sub>	(ii)	d <sup>2</sup> sp <sup>3</sup>
(c) $NO_2^+$	(iii)	sp <sup>3</sup> d
(d) $NH_4^+$	(iv)	sp <sup>3</sup>
	(v)	sp

Choose the **correct** answer from the options given below :

(1) (a)-( i), (b)-( ii), (c)-(v) and (d)-(iii)

(2) (a)-(ii), (b)-(i), (c)-(iv) and (d)-(v)

(3) (a)-(iii), (b)-(i), (c)-(v) and (d)-(iv)

(4) (a)-(iv), (b)-(iii), (c)-(ii) and (d)-(v)

## Official Ans. by NTA (3)

**10.** When silver nitrate solution is added to potassium iodide solution then the sol produced is :

- (1) AgI / I<sup>-</sup> (2) AgI / Ag<sup>+</sup>
  - (3)  $KI / NO_3^-$  (4)  $AgNO_3 / NO_3^-$

# Official Ans. by NTA (1)

**11.** Which of the following molecules does not show stereo isomerism ?

- (1) 3,4-Dimethylhex-3-ene
- (2) 3-Methylhex-1-ene
- (3) 3-Ethylhex-3-ene
- (4) 4-Methylhex-1-ene

# Official Ans. by NTA (3)

**12.** Given below are the statements about diborane

- (a) Diborane is prepared by the oxidation of NaBH<sub>4</sub> with I<sub>2</sub>
- (b) Each boron atom is in sp<sup>2</sup> hybridized state
- (c) Diborane has one bridged 3 centre-2-electron bond
- (d) Diborane is a planar molecule

The option with **correct** statement(s) is -

- (1) (c) and (d) only
- (2) (a) only
- (3) (c) only
- (4) (a) and (b) only

## Official Ans. by NTA (2)



- 13. Which one of the following group-15 hydride is 17. the strongest reducing agent? (1)  $AsH_3$ (2) BiH<sub>3</sub> (4) SbH<sub>3</sub>  $(3) PH_3$ **Official Ans. by NTA (2)** 14. Match List-I with List-II: List-I List-II (a) Chloroprene (b) Neoprene (c) Acrylonitrile (iv) CH<sub>2</sub>=CH-CN (d) Isoprene Choose the correct answer from the options given below : (1) (a) - (iii), (b)-(iv), (c) -(ii), (d) -(i) (2) (a) - (ii), (b)-(iii), (c) -(iv), (d) -(i) (3) (a) - (ii), (b)-(i), (c) -(iv), (d) -(iii) (4) (a) - (iii), (b)-(i), (c) -(iv), (d) -(ii) Official Ans. by NTA (2) 15. The set having ions which are coloured and paramagnetic both is -(1)  $Cu^{2+}$ ,  $Cr^{3+}$ ,  $Sc^{+}$ (2)  $Cu^{2+}$ ,  $Zn^{2+}$ ,  $Mn^{4+}$ (3)  $Sc^{3+}$ ,  $V^{5+}$ ,  $Ti^{4+}$ (4)  $Ni^{2+}$ ,  $Mn^{7+}$ ,  $Hg^{2+}$ Official Ans. by NTA (1) Thiamine and pyridoxine 16. are also known respectively as : (1) Vitamin  $B_2$  and Vitamin E (2) Vitamin E and Vitamin  $B_2$ (3) Vitamin B<sub>6</sub> and Vitamin B<sub>2</sub> (4) Vitamin  $B_1$  and Vitamin  $B_6$ Official Ans. by NTA (4)
  - 7. Sulphide ion is soft base and its ores are common for metals.

(a) Pb	(b) Al
(c) Ag	(d) Mg


Choose the **correct** answer from the options given below :


- (1) (a) and (c) only
- (2) (a) and (d) only
- (3) (a) and (b) only
- (4) (c) and (d) only


## Official Ans. by NTA (1)

18. An organic compound A (C<sub>6</sub>H<sub>6</sub>O) gives dark green colouration with ferric chloride. On treatment with CHCl<sub>3</sub> and KOH, followed by acidification gives compound B. Compound B can also be obtained from compound C on reaction with pyridinium chlorochromate (PCC). Identify A, B and C.

(2) 
$$A = \bigcirc B = \bigcirc CH_2OH OH CHO$$









**19.** Which one of the following reactions does not occur ?

(1) 
$$(1)$$
  $(CH_3CO)_2O/Pyridine \rightarrow (CH_3CO)_2O/Pyridine \rightarrow (CH_3CO)_2$ 

(2) 
$$\bigvee_{SO_3H}^{NH_2} + H_2SO_4 \rightarrow \bigotimes_{SO_3H}^{NH_2}$$

$$(3) \bigoplus^{\mathrm{NH}_2} + \mathrm{AlCl}_3 + \mathrm{CH}_3\mathrm{Cl} \rightarrow \bigoplus^{\mathrm{NH}_2}_{\mathrm{CH}_3}$$

(4) 
$$\bigvee_{NO_2}^{NH_2} + HNO_3/H_2SO_4 \rightarrow \bigotimes_{NO_2}^{NH_2}$$

**Official Ans. by NTA (3)** 

**20.** Which one of the following 0.06 M aqueous solutions has lowest freezing point ?

(1)  $Al_2(SO_4)_3$  (2)  $C_6H_{12}O_6$ (3) KI (4)  $K_2SO_4$ 

Official Ans. by NTA (1)

#### **SECTION-B**

1. The total number of unpaired electrons present in  $[Co(NH_3)_6]Cl_2$  and  $[Co(NH_3)_6]Cl_3$  is

Official Ans. by NTA (1)

ALLEN Ans. (3)

Methylation of 10 g of benzene gave 9.2 g of toluene. Calculate the percentage yield of toluene
 \_\_\_\_\_. (Nearest integer)

# Official Ans. by NTA (78)

**3.** The number of acyclic structural isomers (including geometrical isomers) for pentene are \_\_\_\_\_

# Official Ans. by NTA (6)

4. Assume a cell with the following reaction  $Cu_{(s)} + 2Ag^{+}(1 \times 10^{-3} \text{ M}) \rightarrow Cu^{2+}(0.250 \text{ M}) + 2Ag_{(s)}$   $E^{\Theta}_{cell} = 2.97 \text{ V}$  $E_{cell}$  for the above reaction is V.

(Nearest integer)

[Given :  $\log 2.5 = 0.3979$ , T = 298 K]

Official Ans. by NTA (3)

5. Value of K<sub>P</sub> for the equilibrium reaction

 $N_2O_{4 (g)} \rightleftharpoons 2NO_{2(g)}$  at 288 K is 47.9. The K<sub>C</sub> for this reaction at same temperature is \_\_\_\_\_. (Nearest integer)  $(R = 0.083 \text{ L bar K}^{-1} \text{ mol}^{-1})$ 

Official Ans. by NTA (2)

6. If the standard molar enthalpy change for combustion of graphite powder is  $-2.48 \times 10^2$  kJ mol<sup>-1</sup>, the amount of heat generated on combustion of 1 g of graphite powder is \_\_\_\_\_ kJ. (Nearest integer)

#### Official Ans. by NTA (21)

7. A copper complex crystallising in a CCP lattice with a cell edge of 0.4518 nm has been revealed by employing X-ray diffraction studies. The density of a copper complex is found to be 7.62 g cm<sup>-3</sup>. The molar mass of copper complex is \_\_\_\_\_ g mol<sup>-1</sup>. (Nearest integer)

 $[Given: N_A = 6.022 \times 10^{23} \text{ mol}^{-1}]$ 

## Official Ans. by NTA (106)

 Number of electrons that Vanadium (Z = 23) has in p-orbitals is equal to \_\_\_\_\_

Official Ans. by NTA (12)



9. 
$$N_2O_{5(g)} \rightarrow 2NO_{2(g)} + \frac{1}{2}O_{2(g)}$$

In the above first order reaction the initial concentration of N<sub>2</sub>O<sub>5</sub> is  $2.40 \times 10^{-2}$  mol L<sup>-1</sup> at 318 K. The concentration of N<sub>2</sub>O<sub>5</sub> after 1 hour was 1.60  $\times 10^{-2}$  mol L<sup>-1</sup>. The rate constant of the reaction at 318 K is \_\_\_\_\_\_  $\times 10^{-3}$  min<sup>-1</sup>. (Nearest integer) [Given : log 3 = 0.477, log 5 = 0.699] Official Ans. by NTA (7)

10. If the concentration of glucose  $(C_6H_{12}O_6)$  in blood is 0.72 g L<sup>-1</sup>, the molarity of glucose in blood is  $\_$  × 10<sup>-3</sup>M. (Nearest integer)

[Given : Atomic mass of C = 12, H = 1, O = 16 u]

Official Ans. by NTA (4)