

JEE Main

Shift 2



Physics

- Q.1. An object starts moving with an initial speed 10 m s^{-1} and acceleration 2 m s^{-2} along positive x-direction. The time taken to attain 60 m s^{-1} speed is
- A) 25 s B) 20 s C) 30 s D) 15 s

Answer: 25 s

Solution: The equation of motion for the object is given by

v = u + at ... (1)

Substitute the values of the known parameters into equation (1) and solve to calculate the required time.

 $60 = 10 + 2 \times t$ $\Rightarrow t = \frac{60 - 10}{2}$ = 25

Hence, the required time is $25\ \mathrm{s}$

Q.2. Potential energy of an electron is defined as $U = \frac{1}{2}m\omega^2 x^2$ and follows Bohr's law. Radius of orbit as function of *n* depends on

A)
$$n^2$$
 B) $\frac{1}{\sqrt{n}}$ C) \sqrt{n} D) $n^{2/3}$

Answer: \sqrt{n}

Solution: Here, in the question, *x* is the radius of orbit.

So, we need to find x as a function of n.

i.e., x = f(n)

The angular momentum of an electron in a Bohr orbit is given as below,

$$mvx = \frac{nh}{2\pi}$$
-----(i)

But, $v = x\omega$ -----(ii)

Substituting (ii) in (i), we get

$$m\omega x^2 = \frac{nh}{2\pi}$$

Hence, $x \propto \sqrt{n}$

Q.3. If W is the weight on the surface of Earth then weight of same body at a height $\frac{R_e}{4}$ above the surface of Earth is equal to

 $(R_e = \text{Radius of Earth})$

A)
$$\frac{4}{5}W$$
 B) $\frac{16}{25}W$ C) $\frac{25}{16}W$ D) $\frac{5}{4}W$
Answer: $\frac{16}{25}W$



Solution: The variation of acceleration due to gravity with height from the surface of the Earth can be written as

$$g' = \frac{G_{Me}}{(Re+h)^2} \dots \begin{pmatrix} 1 \end{pmatrix}$$

Substitute $\frac{R_e}{4}$ for *h* into equation (1) and simplify to obtain the new value of the acceleration due to gravity at the given height.

$$g' = \frac{G_M e}{\left(R_e + \frac{R_e}{4}\right)^2}$$
$$= \frac{16G_M e}{25R_e^2}$$
$$= \frac{16}{25}g \dots (2)$$

Multiply m on both sides of equation (2) to obtain the new weight of the object.

$$mg' = rac{16}{25}mg$$

 $\Rightarrow W' = rac{16}{25}W$

Q.4. A proton is projected with speed v in magnetic field B of magnitude 1 T. If angle between velocity and magnetic field is 60° as shown below. Kinetic energy of proton is 2 eV (mass of proton = 1.67×10^{-27} kg, $e = 1.6 \times 10^{-19}$ C). The pitch of the path of proton is approximately



A) $6.28\times 10^{-2}\,\mathrm{m}$ B) $6.28\times 10^{-4}\,\mathrm{m}$

 $6.28 imes10^{-4}\,\mathrm{m}$

C) $3.14 \times 10^{-2} \,\mathrm{m}$

D) $3.14 \times 10^{-4} \text{ m}$

Answer: Solution:



Pitch = $v \cos{(60^{\circ})} \times T$

$$=\sqrt{rac{2KE}{m}} imesrac{1}{2} imesrac{2\pi m}{e_B}$$

$$= \sqrt{2mKE} \times \frac{\pi}{e}$$

$$= \sqrt{2 \times 1.67 \times 10^{-27} \times (2 \times 1.6 \times 10^{-19})} \times \frac{3.14}{1.6 \times 10^{-19}}$$

$$= \sqrt{10.688 \times 10^{-46} \times 1.9625 \times 10^{19}}$$

$$= 3.269 \times 10^{-23} \times 1.9625 \times 10^{19}$$

$$= 6.415 \times 10^{-4} \approx 6.28 \times 10^{-4} \text{ m}$$

Q.5. Find the ratio of root-mean-square speed of oxygen gas molecules to that of hydrogen gas molecules, if temperature of both the gases are same.

Embibe: AI Powered Personalised Adaptive Learning & Outcomes Platform



A)
$$\frac{1}{4}$$
 B) $\frac{1}{16}$ C) $\frac{1}{32}$ D) $\frac{1}{8}$

Answer:

 $\frac{1}{4}$

Solution:

The formula to calculate the rms speed for oxygen gas is given by

$$v_o = \sqrt{\frac{3R_T}{Mo}} \dots \left(1\right)$$

The formula to calculate the rms speed for hydrogen gas is given by

$$v_h = \sqrt{\frac{3RT}{M_h}} \quad \dots \left(2\right)$$

Divide the equations to obtain the ratio of the rms speeds.

$$\frac{v_o}{v_h} = \frac{\sqrt{3RT/Mo}}{\sqrt{3RT/Mh}}$$
$$= \sqrt{\frac{Mh}{Mo}} \dots (3)$$

Substitute the values of the molecular masses of the consecutive gases into equation (3) to calculate the required ratio,

$$\Rightarrow \frac{v_o}{v_h} = \sqrt{\frac{2}{32}}$$
$$= \frac{1}{4}$$

Q.6. In amplitude modulation with carrier frequency (A_c) and modulating frequency (A_m) , modulation index is 60%. If $A_c - A_m = 3$ V, then $A_c + A_m$ is equal to

A) 6 V B) 12 V C) 4 V D) 15 V

Answer: 12 V

Solution: The formula to calculate the modulation index (m) is given by

$$m = \frac{Am}{Ac} \quad \dots \quad \left(1\right)$$

Use the method of Componendo Dividendo on both sides of equation (1) and simplify to obtain the required sum.

$$\frac{m+1}{m-1} = \frac{Am+Ac}{Am-Ac}$$
$$\Rightarrow Am + Ac = \frac{m+1}{m-1}(Am - Ac) \quad \dots \quad \left(2\right)$$

Substitute the values of the known parameters into equation (2) to calculate the required value.

$$egin{array}{lll} A_m + A_c {=} rac{0.6{+}1}{0.6{-}1} imes (-3 \ {
m V}) \ {=} 12 \ {
m V} \end{array}$$

Q.7. An electric dipole is shown in the figure. If it is displaced angularly by a small angle with respect to electric field, then angular frequency of oscillation is given by









Here, dipole oscillates about centre of mass (COM).

Also, COM is towards larger mass and its location is $\frac{l}{3}$ from m.

$$\tau = -qlE\sin\left(\theta\right) = I\alpha$$

For small angles, $\sin(\theta) \approx \theta$,

So,
$$\tau = -qlE\theta = I\alpha$$
---(i)

Calculation of I:

$$I = \frac{m}{2} \left(\frac{2l}{3}\right)^2 + m \left(\frac{l}{3}\right)^2 = \frac{ml^2}{3} - -(ii)$$

Substituting (ii) in (i), we get,

$$\frac{-3ql_E\theta}{ml^2} = \alpha$$

On comparing above equation with,

$$lpha=-\omega^2 heta$$
 We get, $\omega=\sqrt{rac{3qE}{ml}}$

Q.8. In the circuit shown reading of the ideal voltmeter used is equal to _____ volts.







The equivalent resistance (R) of the circuit can be calculated as follows

 $egin{array}{ll} R=3 \,\,\Omega+2 \,\,\Omega \ =5 \,\,\Omega \end{array}$

The current (i) in the circuit can be calculated as follows

$$i = \frac{3 \text{ V}}{5 \Omega}$$
$$= 0.6 \text{ A}$$

Hence, the reading in the voltmeter is given by

$$v = 0.6 \text{ A} \times 2 \Omega$$

= 1.2 V

Q.9. In the given AC circuit, find maximum current through the capacitor



B)

 0.35π A

A) $0.65\pi A$

Answer: 0.65π A

Solution: The formula to calculate the instantaneous charge (q) on the capacitor is given by

 $q = C \mathscr{E}$

 $= 36C\sin(120\pi t) \quad \dots (1)$

The value of the instantaneous current (i) through the circuit is can be calculated as follows

C)

 0.2π A

D)

 0.8π A

$$i = \frac{dq}{dt}$$

$$= \frac{d}{dt} (36C \sin (120\pi t))$$

$$= 4320\pi C \cos (120\pi t) \quad \dots (2)$$

The value of the maximum current is, then, given by

 $i_m = 4320 \pi imes 150 imes 10^{-6} ext{ A} \ pprox 0.65 \pi ext{ A}$

Q.10. Radius of first orbit in H - atom is a_0 . Then de-Broglie wavelength of electron in the third orbit is

A)	$3\pi a_0$	B)	$6\pi a_0$	C)	$9\pi a_0$	D)	$12\pi a_0$
----	------------	----	------------	----	------------	----	-------------

Answer: $6\pi a_0$



Solution: Radius of nth orbit is given by,

$$r = a_0 \frac{n^2}{Z} = a_0 n^2 \left(for \ H - atom, \ Z = 1 \right)$$
 ---(i)

The de-Broglie wavelength of electron is given by,

$$\lambda = \frac{h}{mv}$$
 ---(ii)

We also know by Bohr's law, that

$$mvr = \frac{nh}{2\pi}$$
$$\Rightarrow mv = \frac{nh}{2\pi r} \quad ---(iii)$$

Substituting, (iii) in (ii), we get,

$$\lambda = \frac{2\pi r}{r}$$
 ----(iv)

Substituting (i) in (iv), we get,

$$\lambda = 2\pi a_0 n$$

Hence, de-Broglie wavelength of electron in the third orbit is $6\pi a_0$.

- Q.11. Choose the incorrect statement from the given statements
- A) Planets revolve around the sun with constant linear speed.
- B) Energy of planet in elliptical orbit is constant.
- C) Satellite in circular motion have constant energy.
- D) Body falling towards the earth results in negligible displacement of earth
- Answer: Planets revolve around the sun with constant linear speed.
- Solution: From Kepler's second law of planetary motion, the linear speed of a planet is maximum when its distance from the sun is least. So it changes depending on the position of the planet.

Planets follow two conservation laws: total energy stays constant & angular momentum stays constant throughout the elliptical orbital motion.

In circular motion about the Earth, a satellite remains at a fixed distance from the surface of the Earth at all the time, therefore Satellite in circular motion have constant energy.

As mass of earth is very large compared to any object, during free fall of the object, centre of mass of the system remains constant but due to large mass of the earth very little displacement of earth happens.

Q.12. A particle moves from A to B via C with uniform speed $\pi \text{ m s}^{-1}$. Average velocity during the journey is equal to





Solution: Let's consider the following figure-



From the figure, the displacement of the particle on moving from point A to B via C can be calculated as

 $AB = R\sin 60^{\circ} + R\sin 60^{\circ}$ $= 2R\sin 60^{\circ} \dots (1)$

Also, the time taken (t) by the particle to move from A to B is given by

$$t = \frac{AC}{v}$$
$$= \frac{\frac{2\pi R}{3}}{\frac{3}{\pi}} s$$
$$= \frac{2R}{3} s \qquad \dots (2$$

Hence, the average velocity of the particle (v_a) can be calculated as follows

$$v_a = \frac{AB}{t}$$
$$= \frac{2R\sin 60^\circ \text{ m}}{2\frac{R}{3} \text{ s}}$$
$$= \frac{3\sqrt{3}}{2} \text{ m s}^{-1}$$

Q.13. A solid sphere and a ring have equal masses and equal radius of gyration. If sphere is rotating about its diameter and ring about an axis passing through the centre and perpendicular to its plane, then the ratio of radius is $\sqrt{\frac{x}{2}}$, then find the value of *x*.

A

5

Solution: The formula to calculate the moment of inertia of the sphere is given by

$$I_s = mK^2 = \frac{2}{5}mR_s^2 \dots (1)$$

The formula to calculate the moment of inertia of the ring is given by

$$I_r = mK^2 = mR_r^2 \quad \dots (2)$$

Equate equations (1) and (2) and solve to obtain the required ratio.

$$\frac{\frac{2}{5}mR_s^2 = mR_r^2}{\Rightarrow \frac{Rs^2}{Rr^2} = \frac{5}{2}}$$
$$\Rightarrow \frac{Rs}{Rr} = \sqrt{\frac{5}{2}}$$

Comparing the above equation with the given expression it can be concluded that x = 5.

Chemistry

Q.14. Which of the following compound is most acidic?







B)



C)



D)





Answer:



The acidity of a compound is directly proportional to the stability of its conjugate base. When an acidic compound donates a Solution: proton, it forms a conjugate base. The stability of the conjugate base determines the strength of the acid. Here conjugate base is stable when we have electron withdrawing group (NO_2) by -I effect. The electron-withdrawing effect of the nitro group in meta-nitrophenol increases its acidity by enhancing the ability of the hydroxyl group to donate a proton.

So Compound



is stable among the given.

Q.15.		Nessler's reagent does not	have :				
A)	K	B)	Hg	C)	Ν	D)	I
Answe	er:	Ν					

Answer:

Solution: An alkaline solution of K_2HgI_4 is called Nessler's reagent. Nessler's reagent is used for the qualitative analysis of ammonia. A brown colour precipitate is formed when Nessler's reagent is added to concentrated ammonium salts.

Nessler's reagent does not contain nitrogen, therefore, option C is correct.

Q.16. Which of the following is not obtained on electrolysis of brine solution ?

A)	NaOH	B)	H_2 gas	C)	Cl_2 gas	D)	Na
----	------	----	-----------	----	---------------------------	----	----

Answer: $\mathbf{N}\mathbf{a}$



Sodium chloride and water ionizes as:

 $\mathrm{NaCl}\left(\mathrm{aq}\right) \rightarrow \mathrm{Na}^{+}\left(\mathrm{aq}\right) + \mathrm{Cl}^{-}\left(\mathrm{aq}\right)$ (almost completely ionized) $\mathrm{H_{2}O}\left(l\right)\rightleftharpoons\mathrm{H^{+}}\left(\mathrm{aq}\right)+\mathrm{OH^{-}}\left(\mathrm{aq}\right) \text{ (only slightly ionized)}$

At cathode: Both ${\rm Na^+}(aq)$ and ${\rm H^+}(aq)$ are present near the cathode. Since the discharge potential of ${\rm H^+}$ ions is lower than that of Na^+ ions, therefore, H^+ ions are discharged in preference to Na^+ ions. Hence, H_2 gas is evolved at the cathode while $\ensuremath{\mathrm{Na^+}}$ ions remain in the solution.

 $2\mathrm{H^+} + 2\mathrm{e^-} \rightarrow \mathrm{H}_2(\mathrm{g})$

At anode: Both Cl⁻ and OH⁻ ions are present near the anode. Since the discharge potential of Cl⁻ ions is lower than that of OH^- ions, therefore, CI^- ions are discharged in preference to OH^- ions. Hence, CI_2 gas is evolved at the anode while OH- ions remain in the solution.

 $2 \operatorname{Cl}^-
ightarrow \operatorname{Cl}_2(\mathrm{g}) + 2\mathrm{e}^-$

Therefore, the products formed are ${\rm H}_2,~{\rm Cl}_2~{\rm and}~{\rm NaOH}.$

Therefore, the correct option is D.

ер

Q.17.	Statemer Statemer	nt-1: Morphine nt-2: Morphine	and r narco	nany of its homologues, what what was not the sometimes referred to the sometimes referred to the source of the so	nen ad d to as	ministered in medicinal dos s opiates, since they are ob	es, rel otainec	ieve pain and produce sleep I from the opium poppy.
A) Bo	oth Statem	ent- 1 and Stat	temer	nt- 2 are correct	B)	Both Statement- 1 and St	ateme	nt- 2 are incorrect
C) St	atement-	1 is correct and	l Stat	ement- 2 is incorrect	D)	Statement- 1 is incorrect	and St	atement- 2 is correct
Answer:	Both S	Statement-1 an	d Sta	tement- 2 are correct				
Solution		Morphine and In poisonous of sometimes ref These analges cancer, and in	many doses erred sics a child	y of its homologues, when these produce stupor, co to as opiates, since they a ure chiefly used for the relia- libirth.	adminia ma, co are obt ef of po	stered in medicinal doses, onvulsions and ultimately d ained from the opium popp ostoperative pain, cardiac p	relieve eath. N by. bain ar	pain and produce sleep. Morphine narcotics are nd pains of terminal
Q.18.	Oxidation	n state of ${ m Mn}$ in	1 KM1	${ m nO}_4$ changes by three unit	s in wh	ich medium ?		
A) St	rongly aci	dic	B)	Strongly Basic	C)	Aqueous neutral	D)	Weakly acidic
Answer:	Aqueo	us neutral						
Solution		$ m KMnO_4$ acts a changes to $ m Mn$	as an n^{2+} a	oxidising agent in the neut nd in strongly basic mediu	ral me m it ch	dium and gets reduced to 1 anges to MnO_4^- .	MnO ₂ ,	in acidic medium it
		The oxidation	state	of ${ m Mn}$ in ${ m KMnO_4}$ is $+7$				
		The oxidation	state	of Mn in MnO_2 is $+4$				
		Here the chan	ge in	oxidation state is 3.				
		Therefore, opt	tion C	is correct.				
Q.19.	Which of	the following is	s mos	t basic?				
A) T1	$_2O_3$	I	B)	$\mathrm{Tl}_2\mathrm{O}$	C)	$\mathrm{Cr}_2\mathrm{O}_3$	D)	B_2O_3
Answer:	$\mathrm{Tl}_{2}\mathrm{O}$							
Solution	Lower . Here	r the oxidation s e T1 has least o	state, xidati	more basic is the oxide. In on state and it will be more	Tl ₂ O; basic	$_3$ the oxidation state of Tl is oxide. ${ m Cr}_2{ m O}_3$ is amphoteric	s +3. lı c in na	n ${ m Tl}_2{ m O}$ the oxidation state is $+1$ ture. ${ m B}_2{ m O}_3$ is an acidic oxide.
Q.20.	Which of	the following h	as hi	ghest hydration energy?				
A) Be	2+	I	B)	Mg^{2+}	C)	Ca^{2+}	D)	Ba^{2+}
Answer:	Be^{2+}							



Solutior	n: Be ²⁺ ion has small size and high charge. Hence, it has high polarising power and can attract several water molecules. it has the highest hydration energy among the given ions. Smaller sized and highly charged metal ions have higher hydration energy. In the alkaline earth metals the hydration energy for charged ions is greater than the large-sized cha ions. Therefore, order of hydration energy in this group is								olecules. Thus, nigher sized charged				
		Be	$e^{2+} > Mg^{2+} > C_{2+}$	$a^{2+} >$	${ m Sr}^{2+} > { m Ba}^{2+}.$								
Q.21.	IL	JPA	C name of the co	mpou	nd $\mathrm{K}_3[\mathrm{Co}(\mathrm{C}_2\mathrm{O}_4)_3]$	is:							
A) P	ota	ssiu	m trioxalatocoba	lt(III)			B)	Potassium t	rioxalatocoba	altate (1	II)		
C) F	Pota	ssiu	m cobalttrioxalat	e (II)			D)	Potassium o	oxalatocobalt	tate (III)		
Answer	:	Pot	tassium trioxalato	ocobal	tate (III)								
Solutior	า:	Ligands that include a numerical prefix in the name use the prefixes bis for 2, tris for 3, or tetrakis for 4 to indicate their number.											
		lf t	he complex ion i	s an a	nion, we drop the	ending of	the m	etal name an	d add –ate.				
		In	the given coordi	nation	compound, Co is	the cobalt	meta	, $\mathrm{C_2O_4^{2-}}$ are	the ligands r	named a	as oxalato).	
		T۲	ne charge on [Co	(C_2O_4)	$]_3]$ is -3 and oxal	ate is havi	ng an	oxidation sta	te of -2 .				
		X X	+3(-2) = -3 = +3										
		He	ence, the IUPAC	name	of the compound	K ₃ [Co(C ₂ C	0 ₄) ₃] is	s Potassium t	rioxalatocoba	altate(I	П).		
Q.22.	۷	Vhic	n one doesn't giv	/es Pb	$^{+2}$ test?								
A) lo	odit	Э		B)	Chromate		C)	Sulfate		D)	Nitrate		
Answer	:	Nitr	ate										
Solutior	n:	Pł wł	${ m b}^{+2}$ test is used t nich are usually c	o iden colored	tify the presence d and can be easi	of lead ion ly identified	s in a d.	solution. Lea	id ions react	with ma	any anion	is to form	insoluble salts,
		loo	dite ions and chr	omate	ions both react w	ith lead ior	ns to f	orm insoluble	e lead salts, s	so they	would giv	/e a positi	ive ${ m Pb}^{+2}$ test.
		Su in	Ilfate ions also re the solution.	eact w	th lead ions to for	rm a white	precip	itate of lead	sulfate (PbS)	O_4), ind	licating th	ie presend	ce of lead ions
		Ho foi	owever, nitrate io rm precipitates w	ns do <i>i</i> ith mo	not react with lea st cations, includi	d ions to foing lead ior	orm ar ns. Th	i insoluble sa erefore, nitra	lt. Nitrate ion te ions would	is are u d not gi	sually sol ve a posi	luble in wa tive Pb ⁺²	ater and do not test.
Q.23.	C	ons	ider the following	J									
	(i). D.D.T (ii) Aldrin (iii) Sodium arsinate (iv) Sodium chlorate											
	H	low r	many of these ar	e pest	icides?								
A) 1				B)	2		C)	3		D)	4		
Answer	:	2											
Solutior	n:	Di als	chlorodiphenyltri so a pesticide tha	chloro at was	ethane (DDT) an commonly used t	d Aldrin are	e inde sects	ed pesticides	s. DDT was w	videly u	sed as ar	1 insecticio	de, aldrin is
		Sc	odium arsenate a	and so	dium chlorate are	indeed he	rbicid	es.Sodium ar	senate was o	once us	ed to cor	ntrol weed	ls.
Q.24.	H	low r	many of the follow	wing h	ave same Relativ	e lowering	vapou	ir pressure?					
	(.	A) 11	M NaCl										
	(3) 1.	5 M AlCl ₃										
	(C) 11	M urea										
	() 21	M Na_2SO_4										
Answer	:	2											



Relative lowering vapour pressure can be calculated by using the formula :

$$\frac{\Delta P}{P^{\circ}} = i. X_{Solute} \equiv i. M$$

i = Van't Hoff's factor

X = mole fraction

M = Molarity

$$\begin{aligned} \text{(A)} \ & \frac{\Delta P}{P^{\circ}} = 1 \times 2 \\ \text{(B)} \ & \frac{\Delta P}{P^{\circ}} = 1 \times 1 = 1 \\ \text{(C)} \ & \frac{\Delta P}{P^{\circ}} = 4 \times 1.5 = 6 \\ \text{(D)} \ & \frac{\Delta P}{P^{\circ}} = 3 \times 2 = 6 \end{aligned}$$

Therefore, C and D have same Relative lowering vapour pressure.

Q.25. Types of unit cells are cubic, tetragonal, orthorhombic, hexagonal, monoclinic, triclinic, and rhombohedral.

How many of them can have BCC unit cell ?

Answer:

3

Solution: Cubic system shows three types of Bravais lattices - Primitive, base centered and face centered.

Tetragonal system shows two types of Bravais lattices - Primitive, body centered.

Orthorhombic system shows four types of Bravais lattices - Primitive, body centered, base centered and face centered.

Hexagonal system shows one type of Bravais lattice which is Primitive. Rhombohedral system shows one type of Bravais lattice which is Primitive. Monoclinic system shows two types of Bravais lattices - Primitive, base centered. Triclinic system shows one type of Bravais lattice which is Primitive.

Therefore, the correct answer is 3.

Mathematics

Q.26.

If the coefficient of x^7 in $\left(lpha x^2 + rac{1}{2eta x}
ight)^{11}$ and x^{-7} in $\left(x + rac{1}{3eta x^2}
ight)^{11}$ are equal, then

A)
$$\alpha^{6}\beta = \frac{2^{5}}{3^{6}}$$
 B) $\alpha^{6}\beta = \frac{2^{6}}{3^{5}}$ C) $\alpha\beta^{6} = \frac{2^{5}}{3^{6}}$ D) $\alpha\beta^{6} = \frac{2^{6}}{3^{5}}$
Answer: $\alpha^{6}\beta = \frac{2^{5}}{3^{6}}$



Solution: Given,

The coefficient of
$$x^7$$
 in $\left(lpha x^2+rac{1}{2eta x}
ight)^{11}$ and x^{-7} in $\left(x+rac{1}{3eta x^2}
ight)^{11}$ are equal,

Now finding the coefficient of x^7 in $\left(lpha x^2+rac{1}{2eta x}
ight)^{11}$ by using general term of binomial we get,

$$T_{r+1} = {}^{11}C_r \left(\alpha x^2\right)^{11-r} \left(\frac{1}{2\beta x}\right)^r$$
$$\Rightarrow T_{r+1} = {}^{11}C_r(\alpha)^{11-r} \left(\frac{1}{2\beta}\right)^r x^{22-2r-r}$$

Now solving $22 - 3r = 7 \Rightarrow r = 5$

Hence,
$$T_6 = {}^{11}C_5(\alpha)^6 {\left(\frac{1}{2\beta}\right)}^5 x^7$$

Now finding coefficient of x^{-7} in $\left(x+rac{1}{3eta x^2}
ight)^{11}$ by using general term of binomial expansion we get,

$$\begin{split} T_{r+1} &= {}^{11}C_r(x)^{11-r} {\left(\frac{1}{3\beta x^2}\right)}^r \\ &\Rightarrow T_{r+1} &= {}^{11}C_r(x)^{11-3r} {\left(\frac{1}{3\beta}\right)}^r \end{split}$$

Now equating $11 - 3r = -7 \Rightarrow r = 6$

So,
$$T_7 = {}^{11}C_6(x)^{-7} \left(\frac{1}{3\beta}\right)^6$$

Now equating coefficient of $x^7 \& x^{-7}$ we get,

$$\begin{split} ^{11}C_5(\alpha)^6 \bigg(\frac{1}{2\beta}\bigg)^5 &= {}^{11}C_6\bigg(\frac{1}{3\beta}\bigg)^6 \\ \Rightarrow (\alpha)^6 \bigg(\frac{1}{2\beta}\bigg)^5 &= \bigg(\frac{1}{3\beta}\bigg)^6 \\ \Rightarrow \alpha^6\beta &= \frac{2^5}{3^6} \end{split}$$

Q.27. The system of equations

$$\begin{aligned} x+y+z &= 6\\ x+2y+\alpha z &= 5\\ x+2y+6z &= \beta \end{aligned}$$

- A) Infinitely many solutions for $\alpha = 6$, $\beta = 3$
- C) Unique solutions for $\alpha = 6, \ \beta = 5$

- B) Infinitely many solutions for $\alpha = 6$, $\beta = 5$
- D) No solution for $\alpha = 6$, $\beta = 5$
- Answer: Infinitely many solutions for $\alpha = 6$, $\beta = 5$



Solution: Given,

x+y+z=6

 $x+2y+\alpha z=5$

 $x + 2y + 6z = \beta$

The system of equation can be written as

$$\Delta = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & \alpha \\ 1 & 2 & 6 \end{bmatrix}$$

For Unique solution, $\Delta \neq 0$

 $\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & \alpha \\ 1 & 2 & 6 \end{vmatrix} = (12 - 2\alpha) - (6 - \alpha) + (2 - 2)$

 $= 6 - \alpha$

That means for $\alpha = 6$, $\Delta = 0$

Now when $\alpha = 6$

$$\Delta_{1} = \begin{vmatrix} 6 & 1 & 1 \\ 5 & 2 & 6 \\ \beta & 2 & 6 \end{vmatrix} = 0 - (30 - 6\beta) + (10 - 2\beta)$$
$$= 4 (\beta - 5)$$
$$\Delta_{2} = \begin{vmatrix} 1 & 6 & 1 \\ 1 & 5 & 6 \\ 1 & \beta & 6 \end{vmatrix} = (30 - 6\beta) - 0 + (\beta - 5)$$
$$= 25 - 5\beta$$
$$\Delta_{3} = \begin{vmatrix} 1 & 1 & 6 \\ 1 & 2 & 5 \\ 1 & 2 & \beta \end{vmatrix} = (2\beta - 10) - (\beta - 5) + 0$$
$$= \beta - 5$$

Clearly, at $\beta = 5, \ \Delta_i = 0 \ {\rm for} \ i=1,2,3$

Therefore, at $\alpha = 6, \beta = 5$ system has infinitely many solution.

Q.28. If
$$f(x) + f(\pi - x) = \pi^2$$
, then $\int_0^{\pi} f(x) \sin x dx =$

A)
$$\pi^2$$
 B) $\frac{\pi^2}{2}$ C) $\frac{\pi^2}{4}$ D) $\frac{\pi^2}{3}$

Answer: π^2





 $I = \int_0^{\pi} f(x) \sin x dx \quad \dots (1)$ Now using the property $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$ we get, \Rightarrow I = $\int_0^{\pi} f(\pi - x) \sin{(\pi - x)} dx$ $\Rightarrow I = \int_0^{\pi} f(\pi - x) \sin x \, dx \quad \dots (2)$ Adding (1) & (2), we get $2I = \int_0^{\pi} [f(x) + f(\pi - x)] \sin x \, dx$ $\Rightarrow 2I = \pi^2 \int_0^\pi \sin x \, dx$ $\Rightarrow 2I = \pi^2 [-\cos x]_0^{\pi}$ $\Rightarrow 2I = 2\pi^2$ $\Rightarrow I = \pi^2$ Area (in sq. units) included between y = f(x) = |x-1| + |x-2| and y = 3 is 8 B) C) $\mathbf{2}$ D)

Answer:

Q.29.

A) 4

Solution: Given:

4

$$egin{aligned} y &= f(x) = |x-1| + |x-2| \ \Rightarrow f(x) &= egin{cases} 3 - 2x \colon x < 1 \ 1; & 1 \leq x < 2 \ 2x - 3; & x \geq 2 \end{aligned}$$

Let us draw diagram of y = f(x) & y = 3 we get,



Required area is the area of trapezium which is

 $=rac{1}{2}\left(1+3
ight) imes 2$

= 4 sq. units

Sum of all values of α for which $\hat{\imath} - 2\hat{\jmath} + 3\hat{k}$, $2\hat{\imath} - 3\hat{\jmath} + 4\hat{k}$, $(\alpha + 1)\hat{\imath} + 2\hat{k}$ and $9\hat{\imath} + (\alpha - 8)\hat{\jmath} + 6\hat{k}$ are coplanar. Q.30.

Answer:

None of these



Solution: Let the given vectors be $\overrightarrow{A} = \hat{i} - 2\hat{j} + 3\hat{k}$, $\overrightarrow{B} = 2\hat{i} - 3\hat{j} + 4\hat{k}$, $\overrightarrow{C} = (\alpha + 1)\hat{i} + 2\hat{k}$ and $\overrightarrow{D} = 9\hat{i} + (\alpha - 8)\hat{j} + 6\hat{k}$ We know that if the vectors are coplanar then $\begin{bmatrix} \overrightarrow{AB} & \overrightarrow{AC} & \overrightarrow{AD} \end{bmatrix} = 0$

$$\overrightarrow{AB} = \left(2\hat{1} - 3\hat{j} + 4\hat{k}\right) - \left(\hat{1} - 2\hat{j} + 3\hat{k}\right) = \hat{i} - \hat{j} + \hat{k}$$

$$\overrightarrow{AC} = \left((\alpha + 1)\hat{1} + 2\hat{k}\right) - \left(\hat{1} - 2\hat{j} + 3\hat{k}\right) = \alpha\hat{i} + 2\hat{j} - \hat{k}$$

$$\overrightarrow{AD} = \left(9\hat{1} + (\alpha - 8)\hat{j} + 6\hat{k}\right) - \left(\hat{1} - 2\hat{j} + 3\hat{k}\right) = 8\hat{i} + (\alpha - 6)\hat{j} + 3k$$

Now,

$$\Rightarrow \begin{vmatrix} 1 & -1 & 1 \\ \alpha & 2 & -1 \\ 8 & \alpha - 6 & 3 \end{vmatrix} = 0$$
$$\Rightarrow 1 (6 + \alpha - 6) + (3\alpha + 8) + (\alpha^2 - 6\alpha - 16) = 0$$

On Simplifying we get,

 $\Rightarrow \alpha^2 - 2\alpha - 8 = 0$

Therefore, sum of the roots is -(-2) = 2.

Q.31. If *V* is volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , then volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , the volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , the volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , \vec{b} , \vec{c} , the volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , \vec{b} , \vec{c} , the volume of parallelepiped whose edges determined by vectors \vec{a} , \vec{b} , \vec{c} , \vec{c} , \vec{b} , \vec{c} , \vec{c} , \vec{b} , \vec{c}

A) 6V B) V C) 2V D) 3V

Answer: 3V

Solution: We know that,

Volume of parallelepiped whose edges determined by vectors \overrightarrow{a} , \overrightarrow{b} , \overrightarrow{c} is $\begin{bmatrix} \overrightarrow{a} & \overrightarrow{b} & \overrightarrow{c} \end{bmatrix}$.

Now it is given that V is volume of parallelepiped \vec{a} , \vec{b} , \vec{c} , so required volume of parallelepiped with new edges will be

$$= \begin{bmatrix} \overrightarrow{a} & \overrightarrow{a} + \overrightarrow{b} & \overrightarrow{a} + 2\overrightarrow{b} + 3\overrightarrow{c} \\ = \begin{vmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 2 & 3 \end{vmatrix} \begin{bmatrix} \overrightarrow{a} & \overrightarrow{b} & \overrightarrow{c} \end{bmatrix}$$
$$= 3 \begin{bmatrix} \overrightarrow{a} & \overrightarrow{b} & \overrightarrow{c} \end{bmatrix}$$
$$= 3V$$

Q.32. Consider

 $S_1 \colon (p \Rightarrow q) \lor (\sspace{-}p \land q)$ is a tautology

 $S_2 \colon (q \Rightarrow p) \lor (\ensuremath{\sc r} p \land q)$ is a contradiction

- A) S_1 is true and S_2 is false
- C) Both S_1 and S_2 are false

- B) S_1 is false and S_2 is true
- D) Both S_1 and S_2 are true

Answer: Both S_1 and S_2 are false

Solution:						
Colution.	p	q	p	$r^p \wedge q$	$p \Rightarrow q$	$ q \Rightarrow$
		i H			1	

p	q	$\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $	$\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $	$p \Rightarrow q$	$q {\Rightarrow} p$	$(p \Rightarrow q) \lor (\ \ \ p \land q)$	$(q \Rightarrow p) \lor (\ \ p \land q)$
Т	Т	F	F	Т	Т	Т	Т
Т	F	F	F	F	Т	F	Т
F	Т	Т	Т	Т	F	Т	Т
F	F	Т	F	Т	Т	Т	Т

Hence, $(q \Rightarrow p) \lor (\neg p \land q)$ is a tautology, so both statements are false.

Q.33. If $(21)^{18} + 20(21)^{17} + (20)^2(21)^{16} + \dots + 20^{18} = k \left(21^{19} - 20^{19} \right)$ then $k = 10^{10}$



A)
$$\frac{19}{20}$$
 B) 1 C) $\frac{21}{20}$ D) $\frac{20}{21}$

Answer:

1

Solution:

The given series is
$$(21)^{18} + 20(21)^{17} + (20)^2(21)^{16} + \dots + 20^{18} = k(21^{19} - 20^{19})$$

The given series is in Geometric Progression.

The total number of terms are n = 19.

The first term is $a = 21^{18}$.

And common ratio is $r = \frac{a_2}{a_1} = \frac{20(21)^{17}}{21^{18}} = \frac{20}{21}$

We know that the sum of n terms in GP is given by $S_{\mathrm{n}} = rac{a\left(1-r^n
ight)}{1-r} \; \mathrm{when} \; \; |r| < 1$

$$\Rightarrow S_{19} = \frac{21^{18} \left(1 - \left(\frac{20}{21}\right)^{19}\right)}{1 - \left(\frac{20}{21}\right)}$$
$$= \frac{21^{18} \left(21^{19} - 20^{19}\right)}{21^{19}} \times 21$$
$$= 1 \left(21^{19} - 20^{19}\right)$$

On comparing this with $k\left(21^{19}-20^{19}
ight)$, we get k=1.

Hence, the value of
$$k = 1$$
.

Q.34. If
$$1^2 - 2^2 + 3^2 - 4^2 + \ldots - (2022)^2 + (2023)^2 = m^2 n$$
 where $m, n \in N$ and $m > 19$ then $n - m^2$ is

A) 615 B) 562 C) 812 D) 264

Answer: 615

Solution: Let

$$\begin{split} S &= 1^2 - 2^2 + 3^2 - 4^2 + \ldots + (2021)^2 - (2022)^2 + (2023)^2 \\ \Rightarrow S &= (1-2)(1+2) + (3-4)(3+4) + \ldots + (2021-2022)(2021+2022) + (2023)^2 \\ \Rightarrow S &= -[3+7+11+15+\ldots + 4043] + (2023)^2 \\ \Rightarrow S &= -\frac{1011}{2}(6+1010\times 4) + (2023)^2 \\ \Rightarrow S &= -1011\times 2023 + (2023)^2 \\ \Rightarrow S &= 2023\times 1012 \\ \Rightarrow S &= 34^2 \times 1771 \\ \\ \text{So,} \\ m &= 34, \ n &= 1771 \\ \\ \text{Hence,} \\ n - m^2 &= 615 \\ \\ \\ \text{The rank of the word "PUBLIC" is} \end{split}$$

Answer: 582

Q.35.







Solution: Given that three dice are thrown.

The total number of outcomes when three dice are thrown together is $6^3 = 6 \times 6 \times 6.$

The number of outcomes such that all the outcomes are different is $= 6 \times 5 \times 4$.

For ex: If the outcome in dice 1 is ${}^{\prime\prime}6^{\prime\prime}$ then the number of outcomes for dice 2 should be 5 (1, 2, 3, 4, 5) which excludes the outcome ${}^{\prime\prime}6^{\prime\prime}$.

Hence, the required probability is $=\frac{6{\times}5{\times}4}{6{\times}6{\times}6}$

$$=\frac{20}{36}=\frac{5}{9}=\frac{p}{q}$$

 $\Rightarrow q - p = 9 - 5 = 4$

Therefore, the required answer is 4.