

# **JEE Main**

Shift 1



# **Physics**



#### Q.1. Find the equivalent capacitance across points A and B in the given electrical circuit.



Solution: The variation of acceleration due to gravity with depth can be expressed as follows:

$$g' = g\left(1 - \frac{d}{R}\right) \dots \left(1\right)$$

Substitute  $\frac{R}{2}$  for d into equation (1) to obtain the new acceleration due to gravity at the given depth.

$$g' = g\left(1 - \frac{\frac{R}{2}}{R}\right)$$
$$= \frac{g}{2}$$

Since, the acceleration due to gravity is halved, the weight of the object will also be halved at the given depth.

Hence, the weight of the object at the given depth is 100 N.

Q.4. For an object radiating heat at 300 K, the wavelength corresponding to maximum intensity is  $\lambda$ . If the temperature of body is increased by 300 K, the new wavelength corresponding to maximum intensity will be

A) 
$$\frac{\lambda}{2}$$
 B)  $2\lambda$  C)  $\lambda$  D)  $\frac{5\lambda}{2}$ 

Answer:

 $\frac{\lambda}{2}$ 

Solution: According to Wien's displacement law,

 $\lambda_m T = \text{constant}$ 

where,  $\lambda_m$  is wavelength of maximum intensity and T is absolute temperature.

Let  $\lambda_m$ ' be the required wavelength at 600 K (Since temperature is increased by 300 K)

$$\Rightarrow \lambda_m' imes 600 = \lambda imes 300$$
  
 $\Rightarrow \lambda_m' = rac{\lambda}{2}$ 

Q.5. A particle, when projected at  $15^{\circ}$  with horizontal, has a range of 50 m. Find the range when projected at  $45^{\circ}$  with horizontal.

50 m B) 100 m C) 80 m D) 120 m

Answer: 100 m

Solution:

A)

The formula for horizontal range of a projectile is given by,  $R = rac{u^2 \sin(2 heta)}{g}$ 

For angle of projection,  $\theta = 15^{o}$ , range is given by,

$$R = \frac{u^2 \sin(30^o)}{g}$$
$$= \frac{u^2}{2g} = 50 \text{ m } \left(\text{Given}\right)$$

For angle of projection,  $\theta = 45^{\circ}$ , range is given by,

$$R' = \frac{u^2 \sin(90^o)}{g}$$
$$= \frac{u^2}{g} = 100 \text{ m}$$

Q.6. Statement (1) : An LCR circuit connected to an AC source has maximum average power at resonance.

Statement (2): A resistor only circuit with zero phase difference has maximum average power.

A) (1) and (2) both are correct

B) (1) is correct but (2) is incorrect

C) (1) is incorrect but (2) is correct

D) (1) and (2) both are incorrect

Answer: (1) and (2) both are correct



Solution: We know that at resonance, an LCR circuit connected to an AC source has current amplitude at its maximum value.

In addition, average power reaches a maximum when impedance (*Z*), which depends on the frequency, is a minimum, that is, when  $X_C = X_L$  and Z = R. Thus, at resonance, the average power output of the source in an LCR series circuit is a maximum.

Thus, both the statements are correct.

- Q.7. A monoatomic gas initially at pressure *P* and volume *V* is compressed to  $\frac{1}{8}$  of its volume adiabatically. Final pressure of the gas is equal to
- A) 4P B) 8P C) 16P D) 32P

Answer: 32P

Solution: The initial and final parameters of the gas are related by the equation

$$P_i V_i^{\gamma} = P_f V_f^{\gamma}$$
$$P_f = P_i \left(\frac{V_i}{V_f}\right)^{\gamma} \dots \left(1\right)$$

Substitute the values of the known parameters into equation (1) to obtain the final pressure of the gas.

$$P_{f} = P\left(\frac{V_{i}}{\frac{V_{i}}{8}}\right)^{\frac{5}{3}}$$
$$= 32P$$

Q.8. What is the maximum percentage error in the measurement of quantity *l*, if it is given by  $l = \frac{a^2b^3}{c\sqrt{d}}$ ? Given the percentage error in the calculation of *a*, *b*, *c* and *d* are 1%, 2%, 3% and 4% respectively.

A) 4% B) 12% C) 9% D) 13%

Answer: 13%

Solution: The formula to calculate the percentage error in measuring the quantity *l* can be expressed as

$$\frac{\Delta l}{l}\% = \left[ \left( 2\frac{\Delta a}{a} + 3\frac{\Delta b}{b} + \frac{\Delta c}{c} + \frac{1}{2}\frac{\Delta d}{d} \right) \times 100 \right]\% \quad \dots \left( 1 \frac{\Delta a}{b} + \frac{\Delta c}{b} + \frac{1}{2}\frac{\Delta d}{d} \right) \times 100 \right]\%$$

Substitute the values of the known parameters into equation (1) to calculate the required percentage error to measure the given quantity.

$$\frac{\Delta l}{l} \% = \left[ \left( 2 \times 0.01 + 3 \times 0.02 + 0.03 + \frac{1}{2} \times 0.04 \right) \times 100 \right] \%$$
  
= 13%

Q.9. The equation of progressive wave is  $y = 5 \sin(6t + 0.03x)$ . Find the speed of wave.

Answer: 200

Solution: Here, the direction of wave movement is along -X axis.

The standard equation of a wave moving in -X direction is given by,

 $y = A \sin \left( \omega t + k x \right)$ 

where, the speed of wave is given by, speed =  $\frac{\omega}{k}$ 

Upon substitution, we get

speed = 
$$\frac{6}{0.03}$$
$$= 200 \text{ m s}^{-1}$$

Q.10. Earth shrinks to  $\frac{1}{64}$  times of its initial volume. Time period of Earth rotation is found to be  $\frac{24}{x}$  hr. Find the value of x.



Solution: As the radius of Earth is decreased, its moment of inertia will also change.

The time period (T) and the angular frequency  $(\omega)$  are related by the relation

$$\omega = \frac{2\pi}{T} \dots (1)$$

Considering Earth as a perfect sphere of radius R, its initial angular momentum (L) is given by

$$L = \left(\frac{2}{5}MR^2\right)\omega \quad \dots \left(2\right)$$

From equations (1) and (2), it implies that

$$L = \frac{2}{5}MR^2 \times \frac{2\pi}{T}$$
$$\Rightarrow T = \frac{4\pi}{5L}MR^2 \quad \dots (3)$$

The radius and volume of a spere is related by the relation

$$R \propto V^{\frac{1}{3}} \dots \left(4\right)$$

Using equation (4), the new radius of the Earth (R') is given by

$$\frac{R'}{R} = \left(\frac{\frac{V}{64}}{V}\right)^{\frac{1}{3}}$$
$$\Rightarrow R' = \frac{R}{4} \dots (5)$$

Substitute the expression for the new radius into equation (3) to obtain the new time period (T') when the volume is decreased.

$$T' = \frac{4\pi}{5L} M\left(\frac{R}{4}\right)^2$$
$$= \frac{T}{16} \dots (6)$$

Comparing equation (6) with the given expression for the new time period, it can be concluded that x = 16.

Q.11. A conducting rod of length 1 m is moved across a magnetic field of 0.15 T, with constant speed of  $4 \text{ m s}^{-1}$ . Find force (in N) on rod.

#### Answer:

0

Solution: The formula to calculate the emf ( $\varepsilon$ ) induced in the moving rod of length *l*, moving with a speed of *v* under a magnetic field *B* can be expressed as(in this case length of the rod is perpendicular to both magnetic field and velocity of the rod)

$$\varepsilon = Blv \dots (1)$$

Substitute the values of the known parameters into equation (1) to calculate the required induced emf.

$$arepsilon = 0.15 \mathrm{~T} imes 1 \mathrm{~m} imes 4 \mathrm{~m} \mathrm{~s}^{-1}$$
  
= 0.6 V

Now as the rod is not connected to any circuit, there will not be any current flowing & therefore no magnetic force on the rod.

Hence, the required value of the force is 0.

Q.12. 10 resistors each of 10  $\Omega$  resistance when connected together give minimum equivalent resistance  $R_1$  and maximum equivalent resistance  $R_2$  among various possible combinations. So  $\frac{R_2}{R_1}$  is equal to



A set of resistors connected in a circuit produces maximum value of equivalent resistance when they are connected in series Solution: combination. On the other hand, the equivalent resistance is minimum when the set of resistors are connected in parallel combination.

When ten resistors of resistance 10  $\Omega$  each are connected in series combination, the equivalent resistance ( $R_2$ ) is given by

 $R_2 = 10 \ \Omega + 10 \ \Omega + \dots \dots 10 \ \text{times}$  $= 100 \Omega \ldots (1)$ 

When these are connected in parallel combination, the equivalent resistance  $(R_1)$  of the circuit is given by

 $\frac{1}{R_1} = \frac{1}{10 \Omega} + \frac{1}{10 \Omega} + \dots \dots 10$  times  $\Rightarrow R_1 = 1 \Omega \dots (2)$ 

Divide equation (1) by equation (2) to calculate the required ratio.

 $\frac{R_2}{R_1} = \frac{100 \,\Omega}{1 \,\Omega}$ = 100

Q.13. In an AM wave, amplitude of modulating wave = 3 units and amplitude of carrier wave = 15 units. Find the ratio of maximum to minimum intensity  $\frac{Imax}{I_{min}}$ 

Answer: 2.25

Solution: Given.

Amplitude of modulating wave,  $(A_M) = 3$  units

Amplitude of carrier wave,  $(A_C) = 15$  units

As both modulating and carrier waves are superimposed,  $A_{max} = A_C + A_M$ 

and  $A_{min} = A_C - A_M$ 

Since intensity is proportional to the square of amplitude.

$$\begin{aligned} \frac{Imax}{Imin} &= \left(\frac{AC^{+}AM}{AC^{-}AM}\right)^2 \\ &= \left(\frac{15+3}{15-3}\right)^2 \\ &= \left(\frac{18}{12}\right)^2 \\ &= 1.5^2 = 2.25 \end{aligned}$$

## Chemistry

Q.14.

- Stabiliser used for concentration of sulphide ore is A) Fatty acids B) Pine iol C) Cresols D) **Xanthates**
- Answer: Cresols
- Froth flotation method is used to remove gangue from the sulphide ores by the formation of a powdered ores' suspension in Solution: water with the use of collectors and stabilisers. The role of the stabiliser in the froth flotation process is used to stabilise the froth such as cresols and aniline. It increases the non-wettability of the mineral particles.
- Which of the following complex compound is diamagnetic and low spin? Q.15.

A)	$\left[\mathrm{Co}\left(\mathrm{NH}_{3} ight)_{6} ight]^{+3}$	B)	$[{ m CoCl}_6]^{-3}$	C)	$[CoF_6]^{-3}$	D)	$\left[\mathrm{Fe}\left(\mathrm{H_{2}O}\right)_{6}\right]^{+3}$
Answe	er: $\left[\operatorname{Co}\left(\operatorname{NH}_{3}\right)_{6}\right]^{+3}$						

Solution:  $\ln \left[ \operatorname{Co} \left( \mathrm{NH}_3 \right)_6 \right]^{+3}$  the oxidation state of cobalt is +3. Ammonia is a strong field ligand due to that pairing of electrons takes place. The hybridisation of central metal atom is  $d^2sp^3$ . Since it has no unpaired electrons, it is diamagnetic.

> Strong field ligands cause larger splitting of d orbitals and pairing of electrons is favoured. Hence, it is a low spin complex.



Q.16.	. (	One whic	ch does not st	tabilise	e secondary and tertiar	y protein.				
A)	0 -	O linkaç	ge	B)	$\mathrm{S}-\mathrm{S}$ linkage	C)	Vanderwalls force	D)	Hydrogen bonding	
Answ	er:	0 - 0	linkage							
Soluti	on:	In prir and tl	mary amines, he carboxyl gi	peptid roup o	e linkages (also known f another amino acid, r	as peptide	e bonds) are formed be the formation of a pep	etween the tide chain.	amine group of one amino acid	
		ln seo carbo	condary struct	ture of adjace	polypeptides, the pept ent amino acids.	ide chains	are stabilised by hydro	ogen bond	ing between the amino and	
	In tertiary structure, the protein chain is folded into a three-dimensional conformation, and the stabilised by various types of bonds, including disulphide bonds, ionic bonds, hydrogen bonds, and hydrophobic interactions, among others,									
		So an	nong the give	n optio	ons O – O linkage does	sn't stabilis	e secondary and tertia	ry protein.		
Q.17.	1	Which co	ompound does	s not e	exist from the following?	)				
A)	BeC	$2_1$		B)	$NaO_2$	C)	${ m PbEt}_4$	D)	$(\mathrm{NH}_4)_2\mathrm{BeF}_4$	
Answ	er:	$NaO_2$								
Soluti	Solution: Among the alkali metals, sodium is known to form peroxides, whereas the other alkali metals (such as potassium, rubidium, and cesium) are known to form superoxides. Sodium reacts with oxygen to form sodium oxide ( $Na_2O$ ) as well as with excess oxygen to form sodium peroxide ( $Na_2O_2$ ). However, sodium does not form the superoxide ion ( $O_2^-$ ) under normal conditions. Therefore, the compound that does not exist from the given options is $NaO_2$ .									
Q.18.	F	Prolonga	ated heating o	of ferro	us ammonium sulphate	e is avoide	d to prevent			
A)	Oxic	dation		B)	Reduction	C)	hydrolysis	D)	Breaking	
Answ	er:	Oxidat	ion							
Soluti	ion:	On he ferrou	eating, ferrous us ion oxidised	s sulpł d to fei	nate crystals lose water rric ions.	and anhy	drous ferrous sulphate	$({ m FeSO}_4)$ is	formed. On prolonged heating	
		The r	eaction is as t	follows	5					
		$2\mathrm{FeS}$	$O_4 \rightarrow Fe_2O_3 +$	- SO <sub>2</sub> -	+ SO <sub>3</sub>					
Q.19.	E	Enthalpy	of adsorptior	n and e	enthalpy of formation of	f micelle ai	re respectively			
A)	Pos	itive, Po	sitive	B)	Positive, Negative	C)	Negative, Positive	D)	Negative, Negative	
Answ	er:	Negati	ive, Positive							
Soluti	ion:		The amount adsorbent is Therefore, e	of hea called nthalp	it evolved when one mo enthalpy of adsorption y of adsorption is nega	ble of an a n. In the va ttive.	dsorbate (gas or liquid st majority of cases, ad	) is adsorb dsorption is	ed on the surface of an s an exothermic in nature.	
			Micelle forma means $\Delta H >$	ation d > 0. Th	ecreases the stability o erefore, the enthalpy o	of the collo of formation	idal solution so energy n of micelle is positive.	of the mix	ture should increase which	
Q.20.	F	Read the Statemer Statemer	e following two nt I: Potassiun nt II: $ m K_2 Cr_2 O_7$	o state n dichr 7 is mo	ments. omate is used in volum re soluble in water thar	hetric analy n $\operatorname{Na}_2\operatorname{Cr}_$	/sis. D <sub>7</sub>			
A)	Both	n statem	ents I and II a	re cori	rect.	B)	Both statements I and	d II are inco	prrect.	
C)	Stat	tement I	is correct and	l II is ir	ncorrect.	D)	Statement I is incorre	ect and II is	correct.	
Answ	er:	Staten	nent l is corre	ct and	Il is incorrect.					
Soluti	ion:		Potassium di is hygroscop	ichrom ic in n	ate is preferred over so ature and, therefore, a	odium dich ccurate we	romate in volumetric a eighing is not possible i	nalysis, be n normal a	cause sodium dichromate tmospheric conditions.	
			$\mathrm{Na}_2\mathrm{Cr}_2\mathrm{O}_7$ is	hygro	scopic in nature and is	more solu	ble in water than ${ m K_2Cr}$	<sub>2</sub> O <sub>7</sub> .		
			Therefore, o	ption (	C is correct.					



#### Q.21. Match the following

Column I	Column II
(i). Cotton mills	(a). Biodegradable waste
(ii). Paper mills	(b). Gypsum
(iii). Fertilizers	(c). Non-biodegradable waste
(iv). Thermal power plant	(d). Fly ash

- A) i c, ii a, b; iii c, iv bB) i - a; ii - a, iii - b, iv - d
- $\label{eq:constraint} \textbf{C} \qquad i-a,c; ii-b, iii-b, iv-a \qquad \qquad \textbf{D} \qquad i-c, ii-b,c; iii-b,c, iv-a$
- Answer: i a; ii a, iii b, iv d
- Solution: Cotton mills generate biodegradable waste such as cotton fiber, cottonseed, and other organic materials. Paper mills produce biodegradable waste such as pulp sludge, bark, and wood chips. Fertilizers are made from raw materials such as phosphate rock, sulfur, and potassium salts. During their production, gypsum is generated as a waste product. Thermal power plants generate non-biodegradable waste such as fly ash, bottom ash, and slag as by-products of coal combustion.
- Q.22. Match the column I with column II

Column I	Column II
A. Dacron	P. Thermosetting
B. Urea and formaldehyde resin	Q. Biodegradable
C. Nylon-2, Nylon-6	R. Polyester
D. Nylon-6,6	S. Uses for making bristles of brushes

 $\mathsf{A}) \qquad \mathsf{A}-\mathsf{R}, \mathsf{B}-\mathsf{P}, \mathsf{C}-\mathsf{S}, \mathsf{D}-\mathsf{Q}\mathsf{B}) \qquad \mathsf{A}-\mathsf{P}, \mathsf{B}-\mathsf{R}, \mathsf{C}-\mathsf{Q}, \mathsf{D}-\mathsf{S}\mathsf{C}) \qquad \mathsf{A}-\mathsf{R}, \mathsf{B}-\mathsf{P}, \mathsf{C}-\mathsf{Q}, \mathsf{D}-\mathsf{S}\mathsf{D}) \qquad \mathsf{A}-\mathsf{R}, \mathsf{B}-\mathsf{R}, \mathsf{C}-\mathsf{S}, \mathsf{D}-\mathsf{Q}$ 

Answer: A - R, B - P, C - Q, D - S

Solution: Dacron is a type of polyester fabric made from polyethylene terephthalate (PET). It is a thermoplastic polymer that is commonly used in clothing, bedding, and as a fibrefill material in various products.

Urea formaldehyde resin is a type of thermosetting plastic that is commonly used in the manufacture of adhesives, coatings, and molded products.

Nylon-2 and Nylon-6 are types of biodegradable nylon that can break down naturally in the environment over time. These types of nylon are typically used in applications such as packaging, textiles, and medical products.

Nylon-6, 6 is a type of nylon that is known for its high durability and abrasion resistance. It is commonly used in the manufacture of bristles for brushes, as well as in other applications such as automotive parts, fishing line, and clothing.

Q.23. How many of the following are bent in shape?

 $SO_2, O_3, I_3^-, N_3^-$ .

2



- Solution:  $SO_2$  has a bent molecular geometry due to the presence of a lone pair of electrons on sulfur atom, which distorts the molecular shape. In  $O_3$  central oxygen atom is surrounded by two unshared electrons (one lone pair) and six bonding electrons (three bonds). So it is bent in shape.
  - $I_3^-$  and  $N_3^-$  are linear.



Nitride ion

Q.24. The sum of number of lone pairs in central atom in  $IF_5$  and  $IF_7$  is:

Answer:

1

Solution: In  $IF_5$ , the central atom is iodine, which has 7 valence electrons. Iodine in this molecule has 5 bonding pairs and 1 lone pair. Therefore, the number of lone pairs on the central atom of  $IF_5$  is 1.

In IF<sub>7</sub>, the central atom is also iodine, which has 7 valence electrons. Iodine in this molecule has 7 bond pairs and 0 lone pairs. Therefore, the number of lone pairs on the central atom of IF<sub>7</sub> is 0.

Therefore, the sum of the number of lone pairs in the central atom of IF<sub>5</sub> and IF<sub>7</sub> is 1 + 0 = 1.

Q.25. The degree of dissociation of monobasic acid is 0.3, By what percent is the observed depression in freezing point greater than the calculated depression in freezing point?

Answer: 30

Solution:

 $\overset{\text{HA}}{1-\alpha} \stackrel{\text{H}^+}{=} \overset{\text{H}^+}{\alpha} \overset{\text{A}^-}{+} \overset{\text{A}^-}{\alpha}$ 

Van't Hoff factor,  $i=1-\alpha+\alpha+\alpha=1+\alpha$ 

Given, degree of dissociation,  $\alpha=0.\,3$ 

Therefore, i = 1 + 0.3 = 1.3

The observed depression in freezing point is 1.3 times of the calculated depression in freezing point. Therefore, the difference is 0.3 and the percentage is 30.

Q.26. How many compounds can be easily prepared by Gabriel Phthalimide synthesis, which on reaction with Hinsberg reagent produces a compound which is soluble in KOH?



Answer:

2



Solution: Gabriel's Phthalimide synthesis is used for the conversion of primary alkyl halide into a primary amine. In Gabriel phthalimide synthesis, a base abstract proton from phthalimide gives a nucleophile phthalimide ion which attacks on the unhindered primary alkyl halide. The base hydrolysis of alkylated phthalimide gives the primary unhindered amine and phthalimide ion.



# **Mathematics**

Q.27.	The negation of the st	ateme	nt $(p \lor q) \land  extsf{-}r$ is					
A) (~1	$p \wedge \neg q) \wedge r$	B)	$(\ensuremath{\sc r} p \wedge \ensuremath{\sc -} q) ee r$	C)	$(\ensuremath{{}^{\circ}} p \wedge q) ee r$	D)	$(p \wedge {\scriptscriptstyle{{}^{\sim}}} q) \lor r$	
Answer:	$(\ensuremath{\sc r} p \wedge \ensuremath{\sc r} q) ee r$							
Solution:	The negation of the	e state	ment $(p \lor q) \land  extsf{-}r$ is					
	~ $[(p \lor q) \land  extsf{-}r]$							
	$\equiv$ ~ $(p \lor q) \lor r$							
	$[\because \neg (A \land B) \equiv \neg A \lor$	~ <i>B</i> ]						
	$\equiv ({\scriptstyle{{}^{\sim}}} p \wedge {\scriptstyle{{}^{\sim}}} q) \vee r$							
	$[\because \neg (A \lor B) \equiv \neg A \land \neg$	~ <i>B</i> ]						
Q.28.	From a square of side cuboid with open top is	30 cm s	, the squares of side	$x  ext{ cm}$ is cut	off to make cubo	id of maximum vo	blume. The surfa	ace area of

A)	$400 \text{ cm}^2$	B)	$464 \mathrm{cm}^2$	C)	$800 \text{ cm}^2$	D)	$900 \ \mathrm{cm}^2$
	0						

Answer:  $800 \text{ cm}^2$ 



Solution: Given that, the side of square is 30 cm and x cm squares are cut off.

### The required diagram is



Now the dimensions of the cuboid formed will be

l(x) = 30 - 2x, b(x) = 30 - 2x and h(x) = x.

The Volume of the cuboid will be  $V(x) = (30 - 2x)^2(x)$ 

Now to get Maximum value,

$$\Rightarrow \frac{\mathrm{d} V(x)}{\mathrm{d} x} = 0$$
  
$$\Rightarrow 2 (30 - 2x) (-2)x + (30 - 2x)^2 (1) = 0$$

$$\Rightarrow (30-2x)\left(-4x+30-2x
ight)=0$$

On simplifying we get,

 $\Rightarrow x = 15 \ cm, \ 5 \ cm$ 

But x cannot be  $15\,\,{\rm cm}$  as the volume becomes zero.

Hence x = 5 cm.

Now to find the surface area of the cuboid,



Surface area will be  $= (30 - 2x) imes x imes 4 + (30 - 2x)^2$ 

$$=(30-2 imes 5) imes 5 imes 4+(30-2 imes 5)^2$$

 $= 800 \text{ cm}^2.$ 

Therefore, the required surface area will be  $800 \text{ cm}^2$ .

Q.29. Using the number 1, 2, 3, ..., 7, total numbers of 7 digit number which does not contain string 154 or 2367 is, (repetition is not allowed)

A)	4897	B)	4898	C)	4896	D)	4899
----	------	----	------	----	------	----	------



A) $\sqrt{2}$	B) <sub>1</sub> /5	 C)	$4\sqrt{3}$	D)	None of these
Q.31. S	lope of tangent to a curve at a var	iable point is $\frac{x^2+y^2}{2xy}$ and	$y\left(2 ight)=0$ , then $y\left(8 ight)$ is		
	$=96 imesrac{1}{32}=3$				
	$=96 imesrac{\sin\left(rac{\pi}{33} ight)}{2^5\sinrac{\pi}{33}}~\{\mathrm{as}~\sin\left(\pi-lpha ight)=$	$=\sinlpha\}$			
	$=96 imesrac{\sin\left(\pi-rac{\pi}{33} ight)}{2^5\sinrac{\pi}{33}}$				
	$=96\times\frac{\sin\frac{32\pi}{33}}{2^5\sin\frac{\pi}{33}}$				
	$96 \cdot \cos \frac{\pi}{33} \cdot \cos \frac{2\pi}{33} \cdot \cos \frac{4\pi}{33} \dots \dots$	$.\cos\frac{16\pi}{33}$			
	Now using the above formula in	given expression we get,			
	$\cos A \cdot \cos 2A \cdot \cos 2^2 A \cdot \cos 2^3 A$	$\dots \cdot \cos 2^{n-1}A = \frac{\sin 2^n A}{2^n \sin A}$			
	Now we know that,				
	Expression $96 \cdot \cos \frac{\pi}{33} \cdot \cos \frac{2\pi}{33} \cdot \cos \frac{2\pi}{33}$	$\cos \frac{4\pi}{33} \dots \cos \frac{16\pi}{33}$			
Solution:	Given,	4 10			
Answer:	3				
<b>A)</b> 0	<b>B)</b> 1	C)	2	D)	3
Q.30. F	ind the value of $96 \cdot \cos \frac{\pi}{33} \cdot \cos \frac{2\pi}{33}$	$\cdot \cos \frac{4\pi}{33} \dots \cos \frac{16\pi}{33}$			
	= 5040 - 142 = 4898				
	=5040-(120+24-2)				
	So, total number which does not	7 will be $7! - (5! + 4! - 2)$			
	And number which contain both	string 2367 & 154 will be :	2		
	Total numbers which contain stri	ng 2367 will be 4!			
	Now total numbers which contain	string 154 will be 5!	Suton,		
	I he number 1, 2, 3,, 7	bor will be 71 without ron	atition		
Solution:	Given,				

Answer:  $4\sqrt{3}$ 



Solution: Given:  

$$\frac{dy}{dx} = \frac{x^2 + y^2}{2xy}$$
Put  $y = vx \Rightarrow \frac{dy}{dx} = v + x\frac{dv}{dx}$ 
 $v + x\frac{dv}{dx} = \frac{1 + v^2}{2v}$ 
 $\Rightarrow x\frac{dv}{dx} = \frac{1 - v^2}{2v}$ 
 $\Rightarrow \int \left(\frac{2v}{v^2 - 1}\right) dv = -\int \frac{dx}{x}$ 
 $\Rightarrow \log_e |v^2 - 1| = \log_e \left(\frac{C}{x}\right)$ 
 $\Rightarrow \frac{y^2 - x^2}{x^2} = \frac{C}{x}$ 
 $\Rightarrow y^2 - x^2 = Cx$ 
Put  $x = 2$  and  $y = 0$  we get,  
 $0 - 2^2 = 2C \Rightarrow C = -2$ 
 $\Rightarrow y^2 = x^2 - 2x$ 
 $\Rightarrow y(8) = \sqrt{8^2 - 16}$ 
 $\Rightarrow y(8) = \sqrt{48} = 4\sqrt{3}$ 

Q.32. If the order of the matrix A is  $3 \times 3$  and |A| = 2, then the value of  $\left| 3adj \left( |3A|A^2 \right) \right|$  is

A)  $2^{21} \cdot 3^{10}$  B)  $2^{10} \cdot 3^{21}$  C)  $2^{12} \cdot 3^{15}$  D)  $2^{15} \cdot 3^{12}$ 

Answer:  $2^{10} \cdot 3^{21}$ 

Solution: We need to find the value of  $\left|3adj\left(|3A|A^2\right)\right|$ 

We know that  $|kA| = k^n |A|$  where *n* is the order of the matrix and *k* is a constant.

$$\Rightarrow |3A| = 3^{3}(2)$$

$$\Rightarrow |3adj (|3A|A^{2})| = 3^{3} |adj ((3^{3} \cdot 2)A^{2})|$$
We know that  $adj (kA) = k^{n-1}adj (A)$ 

$$= 3^{3} |(3^{3} \cdot 2)^{2}adj (A^{2})|$$
Now we know that  $|adjA| = |A|^{n-1}$ 

$$= 3^{3} ((3^{3} \cdot 2)^{2})^{3} |adj (A^{2})|$$

$$= 3^{3} (3^{3} \cdot 2)^{6} |A^{2}|^{3-1}$$
  
=  $3^{3} (3^{3} \cdot 2)^{6} (2)^{4}$   
=  $2^{10} \cdot 3^{21}$ 

Hence, the required answer is  $2^{10}\cdot 3^{21}$ 

Q.33. Find the number of integral values of x which satisfy the inequality  $x^2 - 10x + 19 < 6$ 

A) 7		B)	10	C)	6	D)	8
Answer:	7						



Solution: Given:  $x^2 - 10x + 19 < 6$  $\Rightarrow x^2 - 10x + 13 < 0$ Now,  $x^2 - 10x + 13 = 0$  $\Rightarrow x = \frac{10 \pm \sqrt{48}}{2}$  $\Rightarrow x = 5 \pm 2\sqrt{3}$ So.  $x^2 - 10x + 13 < 0$  $\Rightarrow \left(x-\left(5+2\sqrt{3}
ight)
ight)\left(x-\left(5-2\sqrt{3}
ight)
ight)<0$  $\Rightarrow x \in \left(5 - 2\sqrt{3}, 5 + 2\sqrt{3}\right)$  $\Rightarrow x \in (1.5, 8.4)$ So, integral values are x = 2, 3, 4, 5, 6, 7, 8 i.e., 7 values. The coefficient of  $x^7$  in  $\left(1-2x+x^3
ight)^{10}$  is Q.34. B) A) 54102080 C) 4080 D) 6234 Answer: 4080 We need to find the coefficient of  $x^7$  in  $\left(1-2x+x^3
ight)^{10}$ . Solution: We know that for  $(x + y + z)^n$ ,  $T_n = rac{n!}{a^{|b|c|}} (x)^a (y)^b (z)^c$  such that a+b+c=nNow for  $\left(1-2x+x^3
ight)^{10}$  $T_n = \frac{10!}{a!b!c!} (1)^a (-2x)^b \left(x^3\right)^c$  $=\frac{10!}{a!b!c!}-2^bx^{b+3c}$ . with a+b+c=10Here we need coefficient of  $x^7$ . Hence the combinations would be which satisfies both a + b + c = 10 and b + 3c = 7. Hence, coefficient of  $x^7$  is  $=\frac{10!}{3!7!0!}(-2)^7 + \frac{10!}{5!4!1!}(-2)^4 + \frac{10!}{7!1!2!}(-2)^1$  $= 120 \times (-128) + 20160 + (-720)$ =4080Therefore, the required value is 4080 If  $a^2 + {(ar)}^2 + {\left( {ar^2} \right)}^2 = 33033, (a,r \in N)$  then the value of  $a + ar + ar^2$  is Q.35.

C)

230

B)

249

A)

Answer:

148

231

D)

231



Given that  $a^2 + (ar)^2 + (ar^2)^2 = 33033, (a, r \in N)$  $\Rightarrow a^2 \left(1+r^2+r^4
ight) = 11^2 imes 273$ On comparing both sides of the equation we get, a = 11 and  $1 + r^2 + r^4 = 273$  $\Rightarrow r^2 + r^4 = 272$  $\Rightarrow r^2 \left(1+r^2
ight) = 16 imes 17$  $\Rightarrow r = 4$ Now  $a + ar + ar^2 = 11 + 11 \times 4 + 11 \times 16$ = 11 + 44 + 176 = 231.Therefore, the required value is 231 Q.36. Shortest distance between the lines  $\frac{x+1}{7} = \frac{y+1}{-6} = \frac{z+1}{1}$  and  $\frac{x-3}{1} = \frac{y-5}{-2} = \frac{z-7}{1}$  is  $2\sqrt{29}$  units C)  $3\sqrt{29}$  units B) A)  $\sqrt{29}$  units  $2\sqrt{29}$  units Answer: Solution: Given:  $\frac{x+1}{7} = \frac{y+1}{-6} = \frac{z+1}{1}$  $\frac{x-3}{1} = \frac{y-5}{-2} = \frac{z-7}{1}$ So.  $\overrightarrow{a}_1 = -\hat{\imath} - \hat{\jmath} - \hat{k}$  $\overrightarrow{a}_2 = 3\hat{i} + 5\hat{j} + 7\hat{k}$ So.  $\overrightarrow{a}_2 - \overrightarrow{a}_1 = 4\hat{i} + 6\hat{j} + 8\hat{k}$ And.  $\overrightarrow{b}_1 \times \overrightarrow{b}_2 = \begin{vmatrix} \hat{\imath} & \hat{\jmath} & \hat{k} \\ 7 & -6 & 1 \\ 1 & -2 & 1 \end{vmatrix} = -4\hat{\imath} - 6\hat{\jmath} - 8\hat{k}$  $\Rightarrow \left|\overrightarrow{b}_{1} \times \overrightarrow{b}_{2}\right| = \sqrt{16 + 36 + 64} = \sqrt{116}$ Shortest distance between the lines  $= \frac{\left|\frac{\left(\overrightarrow{a}_{2} - \overrightarrow{a}_{1}\right) \cdot \left(\overrightarrow{b}_{1} \times \overrightarrow{b}_{2}\right)}{\left|\overrightarrow{b}_{1} \times \overrightarrow{b}_{2}\right|}\right|}$  $= \left|\frac{\left(4\hat{\imath}+6\hat{\jmath}+8\hat{k}\right)\cdot\left(-4\hat{\imath}-6\hat{\jmath}-8\hat{k}\right)}{\sqrt{116}}\right|$  $=\left|\frac{-16-36-64}{\sqrt{116}}\right|=\sqrt{116}=2\sqrt{29}$  units Q.37. If 3, 8, 13, ..... 373 is in AP then sum of terms which is not divisible by 3 is

9525 Answer:

Solution:

D)

 $5\sqrt{29}$  units



Solution: Given sequence is  $3, 8, 13, \dots, 373$ .

Here a = 3, d = 5 and  $a_n = 373$ 

Now  $a_n = a + (n-1)d$ 

 $\Rightarrow 373 = 3 + (n-1)5$ 

 $\Rightarrow n = 75$ 

We know that  $S_n = \frac{n}{2}(a + a_n)$ 

$$\Rightarrow S_{75} = \frac{75}{2} \left(3 + 373\right)$$

 $\Rightarrow S_{75} = 14100$ 

Now let us write the sequence of terms which are divisible by 3.

We get, 3, 18, 33, ..... 363

$$\Rightarrow 363 = 3 + (n-1)15$$

 $\Rightarrow n = 25$ 

Now let us find the sum of terms divisible by 3.

$$\Rightarrow S_{div\;by\;3} = \frac{25}{2} \left(3 + 363\right) = 4575$$

Required sum = Sum of 75 terms - Sum of terms divisible by 3

= 14100 - 4575

= 9525.

Therefore, the required sum is 9525

Q.38. If the coefficient of  $x^7$  in the expansion of  $\left(ax - \frac{1}{bx^2}\right)^{13}$  is equal to the coefficient of  $x^{-5}$  in the expansion of  $\left(ax + \frac{1}{bx^2}\right)^{13}$ , then  $a^4b^4$  is



Solution: The coefficient of  $x^7$  in the expansion of  $\left(ax^2 - \frac{1}{bx}\right)^{13}$  is equal to the coefficient of  $x^{-5}$  in  $\left(ax + \frac{1}{bx^2}\right)^{13}$ .

We know that, the general term  $\mathrm{T}_{r+1}$  in the expansion  $\left(\mathrm{a}+\mathrm{b}\right)^n~$  is

$$\mathrm{T}_{r+1} = {}^{n} \mathrm{C}_{r} \mathrm{a}^{n-r} \mathrm{b}^{r}$$

Applying to 
$$\left(ax - \frac{1}{bx^2}\right)^{13}$$
, we get  
 $T_{r+1} = {}^{13}C_r(ax){}^{13-r}\left(-\frac{1}{bx^2}\right)^r$   
 $\Rightarrow T_{r+1} = (-1)^r \times {}^{13}C_r(a){}^{13-r}(x){}^{13-3r}(b){}^{-r}$ 

Therefore,  $13 - 3r = 7 \Rightarrow r = 2$  for coefficient of  $x^7$ .

Thus,

$$T_3 = {}^{13}C_2 \left(\frac{a^{11}}{b^2}\right)$$

Similarly, applying to  $\left(ax+rac{1}{bx^2}
ight)^{13}$ , we get

$$T_{r+1} = {}^{13}C_r(ax){}^{13-r}\left(\frac{1}{bx^2}\right)^r$$
$$\Rightarrow T_{r+1} = {}^{13}C_r(a){}^{13-r}(x){}^{13-3r}(b){}^{-r}$$

Therefore, 13 - 3r = -5 for coefficient of  $x^{-5}$ 

 $\Rightarrow r = 6$ 

So,

$$T_7 = {}^{13}C_6(a)^7(b)^{-6}$$

Hence, applying the given condition we get

$$\begin{split} ^{13}C_2 \left(\frac{a^{11}}{b^2}\right) &= {}^{13}C_6(a)^7(b)^{-6} \\ \Rightarrow a^4b^4 &= \frac{13}{13}C_6 \\ \Rightarrow a^4b^4 &= \frac{13!}{7!\cdot 6!} \times \frac{2!\cdot 11!}{13!} \\ \Rightarrow a^4b^4 &= \frac{11\times 10\times 9\times 8}{6\times 5\times 4\times 3} \\ \Rightarrow a^4b^4 &= 22 \end{split}$$

Q.39. Two dice are rolled and sum of two digits is N then probability that  $2^N < N!$  is  $\frac{m}{n}$ , where m and n are coprime, then 11m - 3n is





N is the sum of numbers of two dice.

 $\Rightarrow 2 \leq N \leq 12$ 

Let us check the given condition  $2^N < N!$ 

This condition is only true for  $N \ge 4$ 

We need to find the probability for  $N \ge 4$ .

$$\Rightarrow P(N \ge 4) = 1 - P(N = 2) - P(N = 3)$$

$$= 1 - \frac{1}{36} - \frac{1}{36}$$
$$= \frac{11}{12} = \frac{m}{n}$$

Now let us find 11m - 3n.

$$= 11 \times 11 - 3 \times 12 = 85.$$

Hence, the requried answer is 85.